

Anomalous Hall Effect in non-commutative mechanics

P. A. Horváthy *

Laboratoire de Mathématiques et de Physique Théorique
Université de Tours
Parc de Grandmont
F-37200 TOURS (France)

February 6, 2008

Abstract

The anomalous velocity term in the semiclassical model of a Bloch electron deviates the trajectory from the conventional one. When the Berry curvature (alias noncommutative parameter) is a monopole in momentum space, as found recently in some ferromagnetic crystals while observing the anomalous Hall effect, we get a transverse shift, similar to that in the optical Hall effect.

cond-mat/0606472

1 Introduction

The Anomalous Hall Effect (AHE), characterized by the absence of a magnetic field, is observed in some ferromagnetic crystals. While this has been well established experimentally, its explanation is still controversial. One, put forward by Karplus and Luttinger [1] fifty years ago, suggests that the effect is due to an anomalous current.

Many years later, it has been argued [2] that the semiclassical dynamics of a Bloch electron in a crystal should involve a Berry curvature term, Θ .

*e-mail: horvathy@lmpt.univ-tours.fr.

In the n^{th} band the equations of motion read, in an electromagnetic field,

$$\dot{\mathbf{r}} = \frac{\partial \epsilon_n(\mathbf{k})}{\partial \mathbf{k}} - \dot{\mathbf{k}} \times \boldsymbol{\Theta}, \quad (1)$$

$$\dot{\mathbf{k}} = e\mathbf{E} + e\dot{\mathbf{r}} \times \mathbf{B}(\mathbf{r}), \quad (2)$$

where $\mathbf{r} = (x^i)$ and $\mathbf{k} = (k_j)$ denote the electron's intracell position and quasimomentum, respectively; $\epsilon_n(\mathbf{k})$ is the band energy. The relation (1) exhibits an *anomalous velocity term*, $\dot{\mathbf{k}} \times \boldsymbol{\Theta}$, which is the mechanical counterpart of the anomalous current.

The model is distinguished by the non-commutativity of the position coordinates : in the absence of a magnetic field, $\{x^i, x^j\} = \epsilon^{ijn}\Theta_n$ [3, 4]. In the free noncommutative model in 3 space dimensions, $\boldsymbol{\Theta}$ can only be momentum-dependent such that $\partial_{k_i}\Theta^i = 0$ [3].

A remarkable discovery concerns the AHE in the metallic ferromagnet SrRuO₃. Fang et al. [5] found in fact that the experimental data are consistent with $\boldsymbol{\Theta}$ taking the form of a *monopole in momentum space*,

$$\boldsymbol{\Theta} = \theta \frac{\mathbf{k}}{k^3}, \quad (3)$$

$k \neq 0$. (3) is, furthermore, the only possibility consistent with rotational symmetry [3].

Here we propose to study the AHE in the semiclassical framework (as advocated in [6]), with non-commutative parameter (3). For $\mathbf{B} = 0$ and a constant electric field, $\mathbf{E} = \text{const.}$ and assuming a parabolic profile $\epsilon_n(\mathbf{k}) = \mathbf{k}^2/2$, eqn. (2), $\dot{\mathbf{k}} = e\mathbf{E}$, is integrated as $\mathbf{k}(t) = e\mathbf{E}t + \mathbf{k}_0$. The velocity relation (1) becomes in turn

$$\dot{\mathbf{r}} = \mathbf{k}_0 + e\mathbf{E}t + \frac{e\theta E k_0}{k^3} \hat{\mathbf{n}}, \quad (4)$$

where $\hat{\mathbf{n}} = \hat{\mathbf{k}}_0 \times \hat{\mathbf{E}}$ [“hats” denote vectors normalized to unit length]. The component of $\mathbf{k}_0$ parallel to $\mathbf{E}$ has no interest; we can assume therefore that $\mathbf{k}_0$ is perpendicular to the electric field. Writing $\mathbf{r}(t) = x(t)\hat{\mathbf{k}}_0 + y(t)\hat{\mathbf{E}} + z(t)\hat{\mathbf{n}}$, eqn. (4) yields that the component parallel to $\mathbf{k}_0$ moves uniformly, $x(t) = k_0 t$, and its component parallel to the electric field is uniformly accelerating, $y(t) = \frac{1}{2}eEt^2$. (Our choices correspond to choosing time so that the turning point is at $t = 0$.) However, owing to the anomalous term in (1), the particle is also deviated perpendicularly to $\mathbf{k}_0$ and $\mathbf{E}$, namely by

$$z(t) = \frac{\theta}{k_0} \frac{eEt}{\sqrt{k_0^2 + e^2 E^2 t^2}}. \quad (5)$$




Figure 1: *The anomalous velocity term deviates the trajectory from the plane.*

It follows that the trajectory leaves its initial the plane and suffers, between $t = -\infty$ to $t = \infty$, a *finite transverse shift*, namely

$$\Delta z = \frac{2\theta}{k_0}. \quad (6)$$

Most contribution to the shift comes when the momentum is small, i.e., “near the $\mathbf{k}$ -monopole.”

θ becomes a half-integer upon quantization, $\theta = N/2$, and hence (6) is indeed N/k_0 . The constant $k_0 \neq 0$, the minimal possible value of momentum, plays the role of an impact parameter. Let us observe that while (6) does not depend on the field $\mathbf{E}$ or the electric charge e , the limit $eE \rightarrow 0$ is singular. For $eE = 0$, the motion is uniform along a straight line.

The transverse shift is reminiscent of the recently discovered optical Hall effect [7] and can also be derived, just like in the optical case, using the angular momentum. The free expression [3], $\mathbf{J} = \mathbf{r} \times \mathbf{k} - \theta \hat{\mathbf{k}}$, is plainly broken by the electric field to its component parallel to $\mathbf{E}$,

$$J = J_y = z(t)k_0 - \theta \frac{eEt}{\sqrt{k_0^2 + e^2 E^2 t^2}}, \quad (7)$$

whose conservation yields once again the shift (6).

How can the same argument work for a Bloch electron and for light ? The answer relies, for both problems, on having the same “**k**-monopole” contribution, $-\theta\hat{\mathbf{k}}$ in the angular momentum.

Our model is plainly not realistic: what we described is, rather, the deviation of a freely falling non-commutative particle from the classical parabola found by Galileo. Particles in a metal are not free, though, and their uniform acceleration in the direction of **E** should be damped by some mechanism. It is nevertheless remarkable that we obtain qualitative information from such a toy model.

Note added. I am indebted to Dr. S. Murakami for calling my attention to similar work in the context of the spin Hall effect in semiconductors [8]. I would also like to thank Dr. Y. Kats for informing me about the experimental status of the AHE, see [9].

References

- [1] R. Karplus and J. M. Luttinger, *Phys. Rev.* **95**, 1154 (1954).
- [2] M. C. Chang and Q. Niu, *Phys. Rev. Lett.* **75**, 1348 (1995).
- [3] A. Bérard and H. Mohrbach, *Phys. Rev. D* **69**, 127701 (2004) [hep-th/0310167].
- [4] C. Duval et al. *Mod. Phys. Lett. B* **20**, 373 (2006) [cond-mat/0506051].
- [5] Fang et al. *Science* **302**, 92 (2003) [cond-mat/030232].
- [6] T. Jungwirth, Q. Niu, and A. H. MacDonald, *Phys. Rev. Lett.* **90**, 207208 (2002).
- [7] M. Onoda, S. Murakami, N. Nagaosa, *Phys. Rev. Lett.* **93**, 083901 (2004); K. Yu. Bliokh and Yu. P. Bliokh, *Phys. Lett. A* **333**, 181 (2004) [physics/0402110]; A. Bérard and H. Mohrbach, *Phys. Lett. A* **352**, 190; [hep-th/0404165]; C. Duval, Z. Horváth, P. A. Horváthy, [math-ph/0509031], to appear in *Journ. Geom. Phys.*
- [8] S. Murakami, N. Nagaosa, and S.-C. Zhang, *Science* **301**, 1348 (2003) [cond-mat/0308167]
- [9] Y. Kats et al., *Phys. Rev. B* **70**, 180407(R) (2004).